

## 26. STUDIES ON THE BIOLOGICAL CONTROL OF TWO COMMON VECTOR SNAILS OF PUNJAB BY PREDATORY INSECTS

8 species of aquatic insects have been screened for their predatory activity against two species of snails viz. *Limnoea luteola* and *Indoplanorbis exustus*. Larvae of *Hydaticus* sp. are the most efficient predators of *L. luteola* as they consumed 84.3% of the snails exposed to them and the other species in decreasing order ranked as follows, *Laccotrephes ruber* (70%), *Diplonychus rusticum* (43.3%), *Laccotrephes griseus* (40%) and *Lethocerus indicus* (22.2%). *Laccotrephes ruber* is the most efficient predator of *I. exustus* as it consumed 84.3% of the snails and the other species in decreasing order ranked as follows, *Diplonychus rusticum* (60%), *Hydaticus* sp. larvae (52.5%), *Lethocerus indicus* (20%), *Cybister* sp. (18.7%) and *Laccotrephes griseus* (16.6%). *Ranatra elongata* and *Strenolophus* sp. did not feed either on *L. luteola* or on *I. exustus* and *Cybister* sp. did not feed on *L. luteola*.

### INTRODUCTION

The importance of snails cannot be over emphasised as many of them act as the intermediate hosts of several important diseases of man and livestock namely Schistosomiasis, Fascioliasis and Paramphistomiasis and some are serious crop and garden pests. Their control is, therefore, rather essential. Controlling snails through chemical methods involves large scale dissemination of pesticides or molluscicides which in addition to being expensive may prove hazardous to the health of livestock and man. Studies for the development of alternate methods of snail control are, therefore, quite necessary. The need to develop new methods for the control of medically important snails has also been stressed by the U.S. Parasitic Diseases Panel (Anonymous 1971).

Bequaert (1925, 1926) studied the arthropod enemies of molluscs with particular emphasis on the dipterous parasites of snails. Berg (1953, 1961, 1964a, 1964b, 1973) studied the potential of sciomyzid fly larvae for snail-killing and highlighted their importance for the biological control of snails. The role of sciomyzid fly larvae for the biological control of snails was

also stressed by Neff (1964), Neff & Berg (1966), Knutson *et al.* (1967, 1970), Bratt *et al.* (1969), Eckblad (1971), Ferguson *et al.* (1971) and Geckler (1971). The biological control of snails through giant water bug was studied by Somasunderarao (1963), through another belostomatid bug by Voelker (1968) and through *Hydrophilus* beetles by Maillard (1971). Studies on the biological control of snails as such have also been carried out by Chernin *et al.* (1956, 1971) Michelson (1957), Ferguson *et al.* (1956, 1971), Petitjean (1966), Scott (1970), Yasuvaoka (1970) and Muley (1978).

It was, therefore, considered worthwhile to find out the biological control agents from amongst the aquatic insects living in the water bodies inhabited by snails. The present study deals with the assessment of 8 species of aquatic insects for the biological control of two important species of vector snails namely *Indoplanorbis exustus* and *Limnoea luteola*.

### MATERIALS AND METHODS

Eight species of aquatic insects namely *Laccotrephes ruber* Linn., *L. griseus* Guer. and *Ranatra elongata* Fabr. belonging to Nepidae

MISCELLANEOUS NOTES

(Hemiptera); *Diplonychus rusticum* (Fabr.) and *Lethocerus indicus* (Lep. and Serv.) belonging to Belostomatidae (Hemiptera); *Cybister* sp. and *Hydaticus* sp. larvae belonging to Dytiscidae (Coleoptera); and *Sternolophus* sp. belonging to Hydrophilidae (Coleoptera) were used in the experiments and these were collected from the different permanent water bodies of Ludhiana district. Two species of laboratory bred snails i.e. *Indoplanorbis exustus* and *Limnoea luteola* in well established

aquaria were exposed to the attack of the above aquatic insects and their rates of snail consumption were recorded. Their interesting behaviour patterns were also photographed. The observations were made over a period of about two weeks.

RESULTS AND DISCUSSION

Out of the 8 species of predacious insects used in the present study the larvae of *Hydaticus* sp. ranked first for the control of *Limnoea*

TABLE 1

SHOWING THE CONSUMPTION RATE OF *Limnoea luteola* AND *Indoplanorbis exustus* BY AQUATIC INSECTS ..

Sr. No.	Name of insect	Snail species	No. of snails used	No. of snails left unconsumed	Soft mass/shell consumed	Consumption %
1	2	3	4	5	6	7
1.	<i>Laccotrephes griseus</i> Guer.	<i>Limnoea luteola</i>	20	12(8)	Only soft mass consumed	40
		<i>Indoplanorbis exustus</i>	18	15(3)		16.6
2.	<i>L. ruber</i> Linn.	<i>L. luteola</i>	30	9(21)	—do—	70
		<i>I. exustus</i>	32	5(27)		84.3
3.	<i>Ranatra elongata</i> Fabr.	<i>L. luteola</i>	6	6(0)	—do—	0
		<i>I. exustus</i>	8	5(3)*		
4.	<i>Diplonychus rusticum</i> (Fabr.)	<i>L. luteola</i>	30	17(13)	—do—	43.3
		<i>I. exustus</i>	30	12(18)		60
5.	<i>Lethocerus indicus</i> (Lep. & Serv.)	<i>L. luteola</i>	9	7(2)	—do—	22.2
		<i>I. exustus</i>	20	16(4)		20
6.	<i>Cybister</i> sp.	<i>L. luteola</i>	15	15(0)	—do—	0
		<i>I. exustus</i>	16	12(3)		18.7
7.	<i>Sternolophus</i> sp.	<i>L. luteola</i>	20	20(0)	—do—	0
		<i>I. exustus</i>	22	20(2)*		0.09
8.	<i>Hydaticus</i> sp. larvae	<i>L. luteola</i>	70	11(59)	17 shells completely eaten and 50% shells of the rest partly eaten	84.3
		<i>I. exustus</i>	40	19(21)		52.5

N.B. Figures in parenthesis under heading 5 indicate snails consumed.

\* Natural death.



*luteola* as they ate up 84.3% of the snails exposed to their predatory activity. They were found to be voracious feeders of snails as they even ate up the hard shells along with the soft parts. The other species which ranked next in decreasing order were *Laccotrephes ruber*, *Diplonychus rusticum*, *Laccotrephes griseus* and *Lethocerus indicus* as they consumed 70%, 43.3%, 40% and 22.2% of the *L. luteola* snails respectively (Table 1). The other 3 species namely *Ranatra elongata*, *Cybister* sp. and *Sternolophus* sp. did not feed even on a single *L. luteola* snail. *Laccotrephes ruber* has been found to be the most efficient predator of *Indoplanorbis exustus* as it consumed 84.3% of the snails exposed to its predatory activity. The other species namely *Diplonychus rusticum*, *Hydaticus* sp. larvae, *Lethocerus indicus*, *Cybister* sp. and *Laccotrephes griseus* consumed 60%, 52.5%, 20%,

18.7% and 16.6% of the *I. exustus* snails respectively. *Ranatra elongata* and *Sternolophus* sp. did not consume any *I. exustus* snail although 3 out of 8 snails & 2 out of 22 snails exposed to them respectively died a natural death. Somasundararao (1963) also studied the predatory activity of *Sphaerodema rusticum* (now called as *Diplonychus rusticum*) and found that 39 bugs destroyed 309 snails in 5 days which worked out to be 45 snails per month for each bug. He further observed that *Limnoea luteola* snails were preferred over *Indoplanorbis corneus* and the latter over *Limnoea accuminata*.

#### ACKNOWLEDGEMENT

We are grateful to Dr. Kuldip Rai, Zoologist, Zoological Survey of India, Calcutta for the identification of the predatory insects used in the present study.

DEPARTMENT OF VETY. PARASITOLOGY,  
PUNJAB AGRIL. UNIVERSITY,  
LUDHIANA (PB.),  
November 22, 1982.

H. S. BALI  
SAWAI SINGH  
SUNITA SHARMA

#### REFERENCES

- ANONYMOUS (1971): U. S. Parasitic Diseases Panel. In "United States-Japan Co-operative Medical Science Program Five Year Report 1965-1970". pp. 73-82. United States Department of State. Publication 8598.
- BEQUAERT, J. C. (1925): The arthropod enemies of mollusks with description of a new dipterous parasite from Brazil. *J. Parasit.* 11: 201-212.
- (1926): A dipterous parasite of a snail from Brazil with an account of the arthropod enemies of mollusks. In Medical report of the Hamilton Rice Seventh expedition of the Amazon. *Harvard Inst. for trop. Biol. and Med. Contr.* 4: 292-303.
- BERG, C. O. (1953): Sciomyzid larvae (Diptera) that feed on snails. *J. Parasit.* 39: 630-636.
- (1961): Biology of snail — killing Sciomyzidae (Diptera) of North America and Europe. *Verh. XI. Int. Kongr. Ent.* 1: 197-202.
- (1964a): Snail control in Trematode diseases. The possible value of sciomyzid larvae, snail-killing Diptera. In "Advances in Parasitology" (B. Dawes, ed.), Vol. 2, pp. 259-309. Academic Press, London and New York.
- (1964b): Snail-killing sciomyzid flies: biology of the aquatic species. *Verh. int. Verein. theor. angew. Limnol.* 15: 926-932.
- (1973): Biological control of snail borne diseases. A review. *Expl. Parasit.* 33: 318-330.
- BRATT, A. D., KNUTSON, L. V., FOOTE, B. A. & BERG, C. O. (1969): Biology of *Pherbellia* (Diptera: Sciomyzidae). *Med. Cornell Univ. Agric. Exp. Stn.* 404: 1-247.
- CHERNIN, E., MICHELSON, E. H. & AUGUSTINE, D. L. (1956): Studies on the biological control of schistosome bearing snails. *Am. J. trop. Med. Hyg.*

## MISCELLANEOUS NOTES

5: 297-307.

CHERNIN, E. & PERLSTEIN, J. M. (1971): Protection of snails against miracidia of *S. mansoni* by various aquatic invertebrates. *J. Parasit.* 57: 217-219.

ECKBLAD, J. W. (1971): The population ecology of malacophagous larvae of *Sepedon fuscipennis* and three aquatic snail populations (*Lymnoea palustris*: *Physa integra*: *Gyraulus parvus*). Ph.D. thesis. Cornell University, Ithaca N.Y. 108 pp.

FERGUSON, F. F., OLIVER-GONZALEZ, J. & PALMER, J. R. (1958): Potential for biological control of *Australorbis glabratus*, the intermediate host of Puerto Rican schistosomiasis. *Am. J. trop. Med. Hyg.* 7: 491-493.

GECKLER, R. P. (1971): Laboratory studies of predation of snails by larvae of the marsh fly, *Sepedon tenuicornis* (Diptera: Sciomyzidae). *Can. Ent.* 103: 638-649.

KNUTSON, L. V., NEFF, S. E. & BERG, C. O. (1967): Biology of snail-killing flies from Africa and Southern Spain (Sciomyzidae, *Sepedon*). *Parasitology* 57: 487-505.

KNUTSON, L. V., STEPHENSON, J. W. & BERG, C. O. (1970): Bio-Systematic studies of *Salpicella fasciata* (Meigen) a snail-killing fly (Diptera: Sciomyzidae). *Trans. R. ent. Soc. Lond.* 122(3): 81-100.

MAILLARD, Y. P. (1971): La Malacophagei dans le genre *Hydrophilus* Geoffroy (Ins. Coleopteres: Hydrophilidae): son interet dans le controle naturel des hotes intermediaries d helminthiases. *Comptes rendus des Seances de l' Academie des Sciences. Paris* 272: 2235-2238.

MICHELSON, E. H. (1957): Studies on the biological control of schistosome-bearing snails, predators

and parasites of fresh water Mollusca: A review of literature. *Parasitology* 47: 413-426.

MULEY, E. V. (1978): Biological and chemical control of the vector snail *Melania scabra scabra* (Gastropoda: Prosobranchia): *Bull. Zool. Surv. India* 1(i): 1-5.

NEFF, S. E. (1964): Snail-killing sciomyzid flies: application in biological control. *Verh. int. Verein. theor. angew. Limnol.* 15: 933-939.

NEFF, S. E. & BERG, C. O. (1966): Biology and immature stages of malacophagous Diptera of the genus *Sepedon* (Sciomyzidae). *Bulletin Agricultural Experiment Station, Virginia Polytechnic Institute* 566: 1-113.

PETITJEAN, M. (1966): Le controle biologique des mollusques nuisibles. *Annee biol.* 5: 271-295.

SCOTT, J. A. (1970): "Biological regulation of vectors" a special emphasis area. *J. Parasit.* 56: 253.

SOMASUNDARARAO, G. (1963): A preliminary note on the biological control of fresh water snails by aquatic insect. *Indian Vet. J.* 40(i): 50-52.

VOELKER, J. (1968): Untersuchungen zu Ernahrung, Fort pflanzungs-biologie und Entwicklung Von *Limnogeton fieberi* Mayr (Belostomatidae — Hemiptera) als Beitrag zur. Kenntnis Von Natuerlichen Feinden tropischer Surswasser Schnecken. *Ent. Mitt. Zool. St. Inst. Zool. Mus. Hamb.* 3(60): 1-31.

YASUVAOKA, K. (1970): Some recent research on the biology and control of *Oncomelania* snails in Japan. In "Recent Advances in Researches on Filariasis and Schistosomiasis in Japan". (M. Sasa, ed.) pp. 291-303. Univ. of Tokyo. Press, Tokyo; Univ. Park Press, Baltimore and Manchester.

### 27. NECROPHAGOUS HABIT IN THE GIANT AFRICAN SNAIL, *ACHATINA FULICA FULICA* BOWDICK

(With a plate)

On the sunny day of 18th August, 1982 at 6.45 A.M. one of us (RMS) saw a house lizard, *Hemidactylus frenatus* Schlegel dying in a cemented open space near the bath room of his house. The lizard was lying upside down. In the vicinity a number of giant African snails, *Achatina fulica fulica* Bowdick were also present. Out of them one snail came to-

wards the lizard, crawled on it from the head end, came upto the lower jaw and started devouring it within a couple of minutes. Initially the lizard moved its limbs but after 15 minutes it became motionless. This feeding was continued upto 8.10 A.M. When the snail retired, skin and flesh of the lower jaw and the neck of the lizard were found to be eaten